

Table 7.22C GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	118	105	106	119	154
AGE GROUP					
12 to 17	8	7	5	8	6
18 to 25	37	30	22	25	26
26 or Older	111	100	101	117	150
GENDER					
Male	88	87	87	98	130
Female	70	59	55	63	67
PAST YEAR USE					
TOTAL	31	13	31	26	47
AGE GROUP					
12 to 17	5	3	3	5	2
18 to 25	10	10	8	10	9
26 or Older	28	11	30	24	46
GENDER					
Male	29	12	30	24	43
Female	7	4	9	9	19
PAST MONTH USE					
TOTAL	18	9	18	9	29
AGE GROUP					
12 to 17	3	3	1	3	*
18 to 25	4	8	3	3	3
26 or Older	17	*	17	*	29
GENDER					
Male	18	9	17	4	23
Female	*	*	*	8	18

*Low precision, no estimate reported.

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.22D GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.05	0.04	0.04	0.05	0.06
AGE GROUP					
12 to 17	0.03	0.03	0.02	0.03	0.02
18 to 25	0.11	0.09	0.07	0.08	0.08
26 or Older	0.06	0.05	0.05	0.06	0.08
GENDER					
Male	0.07	0.07	0.07	0.08	0.11
Female	0.06	0.05	0.04	0.05	0.05
PAST YEAR USE					
TOTAL	0.01	0.01	0.01	0.01	0.02
AGE GROUP					
12 to 17	0.02	0.01	0.01	0.02	0.01
18 to 25	0.03	0.03	0.03	0.03	0.03
26 or Older	0.01	0.01	0.02	0.01	0.02
GENDER					
Male	0.02	0.01	0.02	0.02	0.03
Female	0.01	0.00	0.01	0.01	0.01
PAST MONTH USE					
TOTAL	0.01	0.00	0.01	0.00	0.01
AGE GROUP					
12 to 17	0.01	0.01	0.00	0.01	*
18 to 25	0.01	0.03	0.01	0.01	0.01
26 or Older	0.01	*	0.01	*	0.01
GENDER					
Male	0.01	0.01	0.01	0.00	0.02
Female	*	*	*	0.01	0.01

*Low precision, no estimate reported.

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.22P GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2010 Versus 2006-2009

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.9693	0.0745	0.0134	0.2184	
AGE GROUP					
12 to 17	0.0975	0.9253	0.8242	0.3658	
18 to 25	0.0000	0.0128	0.8940	0.2463	
26 or Older	0.1252	0.0194	0.0119	0.1301	
GENDER					
Male	0.2292	0.0366	0.0066	0.1354	
Female	0.0653	0.9167	0.6463	0.9934	
PAST YEAR USE					
TOTAL	0.5315	0.0406	0.1848	0.1901	
AGE GROUP					
12 to 17	0.0195	0.4128	0.1872	0.0329	
18 to 25	0.2071	0.3007	0.7482	0.9567	
26 or Older	0.2269	0.0141	0.1715	0.1133	
GENDER					
Male	0.7768	0.1322	0.3878	0.3656	
Female	0.3156	0.0931	0.2033	0.2233	
PAST MONTH USE					
TOTAL	0.2341	0.1342	0.2242	0.0839	
AGE GROUP					
12 to 17	*	*	*	*	
18 to 25	0.9314	0.0892	0.7944	0.7245	
26 or Older	0.1821	*	0.2302	*	
GENDER					
Male	0.6166	0.4826	0.5345	0.1735	
Female	*	*	*	0.2852	

*Low precision, no estimate reported.

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.23C Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
Standard Errors of Numbers in Thousands, 2006-2010**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	165	170	178	216	256
AGE GROUP					
12 to 17	34	37	31	31	37
18 to 25	98	97	94	111	116
26 or Older	112	116	139	161	204
GENDER					
Male	123	124	126	172	197
Female	96	102	117	110	138
PAST YEAR USE					
TOTAL	92	99	101	134	127
AGE GROUP					
12 to 17	27	28	25	25	29
18 to 25	66	69	64	81	84
26 or Older	53	58	71	89	75
GENDER					
Male	62	72	73	106	97
Female	60	62	64	69	66
PAST MONTH USE					
TOTAL	43	46	54	63	73
AGE GROUP					
12 to 17	14	17	13	13	16
18 to 25	32	37	30	36	46
26 or Older	25	15	42	51	49
GENDER					
Male	33	32	41	54	57
Female	27	31	33	35	37

*Low precision, no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.23D Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
Standard Errors of Percentages, 2006-2010**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.07	0.07	0.07	0.09	0.10
AGE GROUP					
12 to 17	0.14	0.15	0.12	0.13	0.15
18 to 25	0.30	0.30	0.28	0.33	0.34
26 or Older	0.06	0.06	0.07	0.08	0.10
GENDER					
Male	0.10	0.10	0.10	0.14	0.16
Female	0.08	0.08	0.09	0.09	0.11
PAST YEAR USE					
TOTAL	0.04	0.04	0.04	0.05	0.05
AGE GROUP					
12 to 17	0.11	0.11	0.10	0.10	0.12
18 to 25	0.20	0.21	0.19	0.24	0.25
26 or Older	0.03	0.03	0.04	0.05	0.04
GENDER					
Male	0.05	0.06	0.06	0.09	0.08
Female	0.05	0.05	0.05	0.05	0.05
PAST MONTH USE					
TOTAL	0.02	0.02	0.02	0.02	0.03
AGE GROUP					
12 to 17	0.05	0.07	0.05	0.05	0.06
18 to 25	0.10	0.11	0.09	0.11	0.13
26 or Older	0.01	0.01	0.02	0.03	0.02
GENDER					
Male	0.03	0.03	0.03	0.04	0.05
Female	0.02	0.02	0.03	0.03	0.03

*Low precision, no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.23P Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
P Values from Tests of Differences of Percentages, 2010 Versus 2006-2009**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.0000	0.0000	0.0000	0.0002	
AGE GROUP					
12 to 17	0.8789	0.7648	0.1862	0.3245	
18 to 25	0.0000	0.0000	0.0000	0.0516	
26 or Older	0.0000	0.0000	0.0000	0.0013	
GENDER					
Male	0.0000	0.0000	0.0000	0.0061	
Female	0.0000	0.0000	0.0000	0.0045	
PAST YEAR USE					
TOTAL	0.0000	0.0000	0.0000	0.4573	
AGE GROUP					
12 to 17	0.9871	0.4163	0.3192	0.5248	
18 to 25	0.0000	0.0000	0.0000	0.1274	
26 or Older	0.0002	0.0014	0.0399	0.6317	
GENDER					
Male	0.0000	0.0002	0.0001	0.9068	
Female	0.0050	0.0000	0.0019	0.2441	
PAST MONTH USE					
TOTAL	0.0000	0.0000	0.0011	0.4432	
AGE GROUP					
12 to 17	0.5312	0.8892	0.2110	0.6554	
18 to 25	0.0083	0.0219	0.0111	0.5747	
26 or Older	0.0005	0.0000	0.0410	0.6641	
GENDER					
Male	0.0004	0.0004	0.0054	0.5602	
Female	0.0175	0.0063	0.0410	0.6128	

*Low precision, no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.24C Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
Standard Errors of Numbers in Thousands, 2006-2010**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	205	215	231	263	289
AGE GROUP					
12 to 17	24	24	21	25	21
18 to 25	61	55	53	55	62
26 or Older	190	206	220	252	279
GENDER					
Male	143	149	149	184	213
Female	140	161	179	178	192
PAST YEAR USE					
TOTAL	138	140	151	149	184
AGE GROUP					
12 to 17	18	16	16	18	15
18 to 25	41	38	34	37	36
26 or Older	130	133	143	140	177
GENDER					
Male	101	98	94	106	143
Female	93	98	112	102	114
PAST MONTH USE					
TOTAL	85	54	58	83	134
AGE GROUP					
12 to 17	10	7	6	7	8
18 to 25	17	17	17	18	16
26 or Older	81	51	55	79	132
GENDER					
Male	70	27	38	49	113
Female	42	45	44	66	73

*Low precision, no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.24D Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
Standard Errors of Percentages, 2006-2010**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.08	0.09	0.09	0.10	0.11
AGE GROUP					
12 to 17	0.10	0.09	0.08	0.10	0.09
18 to 25	0.19	0.17	0.16	0.16	0.18
26 or Older	0.10	0.11	0.11	0.13	0.14
GENDER					
Male	0.12	0.12	0.12	0.15	0.17
Female	0.11	0.13	0.14	0.14	0.15
PAST YEAR USE					
TOTAL	0.06	0.06	0.06	0.06	0.07
AGE GROUP					
12 to 17	0.07	0.06	0.06	0.07	0.06
18 to 25	0.13	0.12	0.10	0.11	0.11
26 or Older	0.07	0.07	0.07	0.07	0.09
GENDER					
Male	0.08	0.08	0.08	0.09	0.12
Female	0.07	0.08	0.09	0.08	0.09
PAST MONTH USE					
TOTAL	0.03	0.02	0.02	0.03	0.05
AGE GROUP					
12 to 17	0.04	0.03	0.02	0.03	0.03
18 to 25	0.05	0.05	0.05	0.05	0.05
26 or Older	0.04	0.03	0.03	0.04	0.07
GENDER					
Male	0.06	0.02	0.03	0.04	0.09
Female	0.03	0.04	0.03	0.05	0.06

*Low precision, no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.24P Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
P Values from Tests of Differences of Percentages, 2010 Versus 2006-2009**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.0000	0.0026	0.0224	0.5986	
AGE GROUP					
12 to 17	0.0250	0.0619	0.3127	0.1102	
18 to 25	0.6404	0.0956	0.4008	0.8388	
26 or Older	0.0000	0.0048	0.0269	0.4792	
GENDER					
Male	0.0001	0.0044	0.0013	0.3121	
Female	0.0005	0.1920	0.9896	0.7554	
PAST YEAR USE					
TOTAL	0.5259	0.5176	0.4654	0.7183	
AGE GROUP					
12 to 17	0.0615	0.2907	0.5969	0.4591	
18 to 25	0.0545	0.3462	0.4071	0.0797	
26 or Older	0.1986	0.3377	0.3255	0.4220	
GENDER					
Male	0.4912	0.4804	0.2181	0.7366	
Female	0.8453	0.8575	0.8337	0.8652	
PAST MONTH USE					
TOTAL	0.1566	0.0045	0.0120	0.2089	
AGE GROUP					
12 to 17	0.4799	0.6245	0.4360	0.9950	
18 to 25	0.7626	0.7293	0.2883	0.3132	
26 or Older	0.1462	0.0041	0.0071	0.1511	
GENDER					
Male	0.4160	0.0092	0.0376	0.2279	
Female	0.1353	0.2229	0.1534	0.6075	

*Low precision, no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.25C Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	282	260	274	257	288
AGE GROUP					
12 to 17	40	41	36	36	39
18 to 25	77	74	71	82	79
26 or Older	263	243	259	235	268
GENDER					
Male	229	202	223	193	236
Female	150	162	169	162	157
PAST YEAR USE					
TOTAL	99	85	98	111	162
AGE GROUP					
12 to 17	30	29	26	25	27
18 to 25	38	37	31	35	36
26 or Older	87	72	91	100	158
GENDER					
Male	76	55	81	75	147
Female	64	66	56	80	70
PAST MONTH USE					
TOTAL	51	46	53	60	67
AGE GROUP					
12 to 17	13	15	13	13	12
18 to 25	15	19	16	17	20
26 or Older	47	39	49	57	63
GENDER					
Male	45	34	43	37	52
Female	26	31	31	47	43

*Low precision, no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.25D Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.11	0.10	0.11	0.10	0.11
AGE GROUP					
12 to 17	0.16	0.16	0.15	0.15	0.16
18 to 25	0.24	0.23	0.22	0.24	0.23
26 or Older	0.14	0.13	0.14	0.12	0.14
GENDER					
Male	0.19	0.17	0.18	0.16	0.19
Female	0.12	0.13	0.13	0.13	0.12
PAST YEAR USE					
TOTAL	0.04	0.03	0.04	0.04	0.06
AGE GROUP					
12 to 17	0.12	0.12	0.10	0.10	0.11
18 to 25	0.12	0.11	0.09	0.10	0.11
26 or Older	0.05	0.04	0.05	0.05	0.08
GENDER					
Male	0.06	0.05	0.07	0.06	0.12
Female	0.05	0.05	0.04	0.06	0.05
PAST MONTH USE					
TOTAL	0.02	0.02	0.02	0.02	0.03
AGE GROUP					
12 to 17	0.05	0.06	0.05	0.05	0.05
18 to 25	0.05	0.06	0.05	0.05	0.06
26 or Older	0.02	0.02	0.03	0.03	0.03
GENDER					
Male	0.04	0.03	0.04	0.03	0.04
Female	0.02	0.02	0.02	0.04	0.03

*Low precision, no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.25P Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2010 Versus 2006-2009

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.9950	0.0338	0.3453	0.1533	
AGE GROUP					
12 to 17	0.0126	0.2083	0.6659	0.7795	
18 to 25	0.4428	0.0943	0.5967	0.7667	
26 or Older	0.8784	0.0613	0.3939	0.1212	
GENDER					
Male	0.9081	0.0197	0.1246	0.0208	
Female	0.8748	0.7507	0.5628	0.5144	
PAST YEAR USE					
TOTAL	0.4752	0.0346	0.0932	0.1375	
AGE GROUP					
12 to 17	0.0152	0.2087	0.4881	0.9259	
18 to 25	0.3763	0.7764	0.9176	0.9306	
26 or Older	0.0935	0.0082	0.0584	0.1134	
GENDER					
Male	0.2970	0.0127	0.0751	0.0278	
Female	0.6812	0.9810	0.8691	0.5128	
PAST MONTH USE					
TOTAL	0.7310	0.5210	0.5033	0.5880	
AGE GROUP					
12 to 17	0.1739	0.1554	0.6082	0.4414	
18 to 25	0.5881	0.7371	0.5048	0.9947	
26 or Older	0.5642	0.2175	0.5263	0.4580	
GENDER					
Male	0.7076	0.5615	0.7251	0.4374	
Female	0.2717	0.7519	0.5317	1.0000	

*Low precision, no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.26C Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	145	140	135	152	166
AGE GROUP					
12 to 17	14	12	12	12	15
18 to 25	50	47	40	37	39
26 or Older	133	129	125	146	160
GENDER					
Male	123	114	107	124	152
Female	69	68	72	78	64
PAST YEAR USE					
TOTAL	38	31	28	32	36
AGE GROUP					
12 to 17	8	7	9	8	11
18 to 25	14	18	13	16	22
26 or Older	34	24	22	27	26
GENDER					
Male	35	27	25	23	33
Female	14	16	11	23	13
PAST MONTH USE					
TOTAL	18	11	7	24	22
AGE GROUP					
12 to 17	4	5	5	6	5
18 to 25	6	6	6	9	8
26 or Older	16	8	*	21	20
GENDER					
Male	14	10	6	13	21
Female	10	4	4	20	8

*Low precision, no estimate reported.

NOTE: Ketamine is also known as "Special K" or "Super K."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.26D Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.06	0.06	0.05	0.06	0.07
AGE GROUP					
12 to 17	0.05	0.05	0.05	0.05	0.06
18 to 25	0.15	0.14	0.12	0.11	0.11
26 or Older	0.07	0.07	0.07	0.08	0.08
GENDER					
Male	0.10	0.10	0.09	0.10	0.12
Female	0.05	0.05	0.06	0.06	0.05
PAST YEAR USE					
TOTAL	0.02	0.01	0.01	0.01	0.01
AGE GROUP					
12 to 17	0.03	0.03	0.04	0.03	0.05
18 to 25	0.04	0.06	0.04	0.05	0.06
26 or Older	0.02	0.01	0.01	0.01	0.01
GENDER					
Male	0.03	0.02	0.02	0.02	0.03
Female	0.01	0.01	0.01	0.02	0.01
PAST MONTH USE					
TOTAL	0.01	0.00	0.00	0.01	0.01
AGE GROUP					
12 to 17	0.02	0.02	0.02	0.03	0.02
18 to 25	0.02	0.02	0.02	0.03	0.02
26 or Older	0.01	0.00	*	0.01	0.01
GENDER					
Male	0.01	0.01	0.00	0.01	0.02
Female	0.01	0.00	0.00	0.02	0.01

*Low precision, no estimate reported.

NOTE: Ketamine is also known as "Special K" or "Super K."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.26P Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2010 Versus 2006-2009

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.8181	0.4307	0.2439	0.7484	
AGE GROUP					
12 to 17	0.9915	0.5227	0.6518	0.8883	
18 to 25	0.0000	0.0000	0.0752	0.3938	
26 or Older	0.0414	0.0405	0.1050	0.6067	
GENDER					
Male	0.2667	0.1517	0.0371	0.3614	
Female	0.0647	0.2424	0.1502	0.2840	
PAST YEAR USE					
TOTAL	0.5019	0.5539	0.0288	0.3774	
AGE GROUP					
12 to 17	0.1497	0.1156	0.2212	0.2503	
18 to 25	0.0297	0.7747	0.0193	0.1233	
26 or Older	0.3673	0.9938	0.6067	0.6766	
GENDER					
Male	0.5495	0.5168	0.0696	0.1304	
Female	0.7456	0.9727	0.1641	0.5019	
PAST MONTH USE					
TOTAL	0.8344	0.4063	0.1516	0.4294	
AGE GROUP					
12 to 17	0.6828	0.7811	0.5762	0.7177	
18 to 25	0.4263	0.6133	0.5648	0.7407	
26 or Older	0.8750	0.5119	*	0.5188	
GENDER					
Male	0.8779	0.6115	0.2955	0.8900	
Female	0.8770	0.3157	0.2182	0.2972	

*Low precision, no estimate reported.

NOTE: Ketamine is also known as "Special K" or "Super K."

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.27C DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	62	71	74	79	101
AGE GROUP					
12 to 17	9	10	10	10	10
18 to 25	29	24	28	33	36
26 or Older	54	65	68	72	94
GENDER					
Male	55	66	60	71	87
Female	28	23	42	32	53
PAST YEAR USE					
TOTAL	21	15	27	29	35
AGE GROUP					
12 to 17	4	6	8	8	8
18 to 25	15	12	14	19	16
26 or Older	14	*	21	20	31
GENDER					
Male	21	14	23	19	28
Female	5	6	14	22	21
PAST MONTH USE					
TOTAL	7	*	18	20	28
AGE GROUP					
12 to 17	*	*	4	4	7
18 to 25	5	4	9	4	7
26 or Older	*	*	15	19	27
GENDER					
Male	6	4	17	8	22
Female	4	*	*	18	18

*Low precision, no estimate reported.

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.27D DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.03	0.03	0.03	0.03	0.04
AGE GROUP					
12 to 17	0.04	0.04	0.04	0.04	0.04
18 to 25	0.09	0.07	0.08	0.10	0.10
26 or Older	0.03	0.03	0.04	0.04	0.05
GENDER					
Male	0.05	0.05	0.05	0.06	0.07
Female	0.02	0.02	0.03	0.02	0.04
PAST YEAR USE					
TOTAL	0.01	0.01	0.01	0.01	0.01
AGE GROUP					
12 to 17	0.02	0.02	0.03	0.03	0.03
18 to 25	0.05	0.04	0.04	0.06	0.05
26 or Older	0.01	*	0.01	0.01	0.02
GENDER					
Male	0.02	0.01	0.02	0.02	0.02
Female	0.00	0.00	0.01	0.02	0.02
PAST MONTH USE					
TOTAL	0.00	*	0.01	0.01	0.01
AGE GROUP					
12 to 17	*	*	0.02	0.02	0.03
18 to 25	0.02	0.01	0.03	0.01	0.02
26 or Older	*	*	0.01	0.01	0.01
GENDER					
Male	0.00	0.00	0.01	0.01	0.02
Female	0.00	*	*	0.01	0.01

*Low precision, no estimate reported.

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.27P DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: P Values from Tests of Differences of Percentages, 2010 Versus 2006-2009

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.0000	0.0000	0.0002	0.0098	
AGE GROUP					
12 to 17	0.2088	0.1930	0.3401	0.8202	
18 to 25	0.0104	0.0003	0.0011	0.1226	
26 or Older	0.0009	0.0037	0.0095	0.0368	
GENDER					
Male	0.0006	0.0017	0.0001	0.0332	
Female	0.0187	0.0033	0.3109	0.1413	
PAST YEAR USE					
TOTAL	0.0030	0.0001	0.0723	0.3862	
AGE GROUP					
12 to 17	0.0170	0.0678	0.4321	0.9894	
18 to 25	0.0310	0.0021	0.0589	0.8143	
26 or Older	0.1098	*	0.4181	0.3687	
GENDER					
Male	0.0399	0.0026	0.1942	0.2045	
Female	0.0167	0.0137	0.1941	0.8943	
PAST MONTH USE					
TOTAL	0.0408	*	0.3089	0.5030	
AGE GROUP					
12 to 17	*	*	0.2212	0.4415	
18 to 25	0.5054	0.1629	0.9211	0.2622	
26 or Older	*	*	0.4422	0.7890	
GENDER					
Male	0.1019	0.0527	0.7486	0.2344	
Female	0.2019	*	*	0.8628	

*Low precision, no estimate reported.

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.28C Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Numbers in Thousands, 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	92	110	127	158	189
AGE GROUP					
12 to 17	20	23	27	28	28
18 to 25	61	68	74	88	95
26 or Older	59	77	97	119	155
GENDER					
Male	80	91	103	141	159
Female	34	54	62	63	99
PAST YEAR USE					
TOTAL	48	68	69	72	77
AGE GROUP					
12 to 17	16	18	23	22	22
18 to 25	38	45	47	47	42
26 or Older	22	43	42	47	59
GENDER					
Male	41	55	55	58	73
Female	21	34	37	37	28
PAST MONTH USE					
TOTAL	18	26	24	28	28
AGE GROUP					
12 to 17	7	9	10	9	9
18 to 25	15	21	21	15	17
26 or Older	*	11	8	22	20
GENDER					
Male	16	22	22	19	27
Female	7	11	8	20	9

*Low precision, no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.28D Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Standard Errors of Percentages, 2006-2010

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.04	0.04	0.05	0.06	0.07
AGE GROUP					
12 to 17	0.08	0.09	0.11	0.12	0.12
18 to 25	0.19	0.21	0.22	0.26	0.28
26 or Older	0.03	0.04	0.05	0.06	0.08
GENDER					
Male	0.07	0.08	0.08	0.12	0.13
Female	0.03	0.04	0.05	0.05	0.08
PAST YEAR USE					
TOTAL	0.02	0.03	0.03	0.03	0.03
AGE GROUP					
12 to 17	0.06	0.07	0.09	0.09	0.09
18 to 25	0.12	0.14	0.14	0.14	0.12
26 or Older	0.01	0.02	0.02	0.02	0.03
GENDER					
Male	0.03	0.05	0.05	0.05	0.06
Female	0.02	0.03	0.03	0.03	0.02
PAST MONTH USE					
TOTAL	0.01	0.01	0.01	0.01	0.01
AGE GROUP					
12 to 17	0.03	0.03	0.04	0.04	0.04
18 to 25	0.05	0.06	0.06	0.04	0.05
26 or Older	*	0.01	0.00	0.01	0.01
GENDER					
Male	0.01	0.02	0.02	0.02	0.02
Female	0.01	0.01	0.01	0.02	0.01

*Low precision, no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

Table 7.28P Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: *P* Values from Tests of Differences of Percentages, 2010 Versus 2006-2009

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010
LIFETIME USE					
TOTAL	0.0000	0.0000	0.0000	0.0009	
AGE GROUP					
12 to 17	0.0000	0.0020	0.7222	0.7780	
18 to 25	0.0000	0.0000	0.0000	0.0421	
26 or Older	0.0000	0.0000	0.0000	0.0054	
GENDER					
Male	0.0000	0.0000	0.0000	0.0164	
Female	0.0000	0.0000	0.0004	0.0095	
PAST YEAR USE					
TOTAL	0.0000	0.1244	0.6859	0.3150	
AGE GROUP					
12 to 17	0.0009	0.1177	0.3690	0.7321	
18 to 25	0.2049	0.5609	0.0664	0.0210	
26 or Older	0.0001	0.0402	0.1853	0.5338	
GENDER					
Male	0.0003	0.0617	0.7148	0.6598	
Female	0.0065	0.7428	0.0958	0.1634	
PAST MONTH USE					
TOTAL	0.4868	0.2554	0.2721	0.3998	
AGE GROUP					
12 to 17	0.1475	0.7839	0.3053	0.6864	
18 to 25	0.5747	0.0907	0.1128	0.8807	
26 or Older	*	1.0000	0.4964	0.4125	
GENDER					
Male	0.5390	0.5274	0.3657	0.9876	
Female	0.7331	0.1241	0.3775	0.1358	

*Low precision, no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.